

EX/SC/MATH/UG/MAJOR/TH/21/205/2025

**B. SC. MATHEMATICS EXAMINATION, 2025**

**( 2<sup>nd</sup> Year, 1<sup>st</sup> Semester )**

**MATHEMATICS**

**PAPER : MAJOR/205**

**( Riemann Integration and Series of Functions )**

*Time : Two Hours*

*Full Marks : 40*

*Use separate Answer-Scripts for each Part.*

*Symbols and Notations have their usual meanings.*

**PART—I (20 Marks)**

Answer *any three* questions from **1 to 5** and **one** question from **6 to 7**.

1. Let a function  $f:[a,b] \rightarrow \mathbb{R}$  be bounded on  $[a, b]$ . Then show that  $f$  is  $R$ -integrable on  $[a, b]$  iff for each  $\varepsilon > 0 \exists \delta(>0)$  such that  $U(p, f) - L(p, f) < \varepsilon$  for every partition  $P$  of  $[a, b]$  with  $\|p\| \geq \delta$ . 5 [CO1]
2. Let  $f:[a,b] \rightarrow \mathbb{R}$  be  $R$ -integrable on  $[a, b]$ . If there exists a  $R \in \mathbb{R}^+$  such that  $f(x) \geq R \forall x \in [a, b]$ , then prove that  $\frac{1}{f}$  is  $R$ -integrable on  $[a, b]$ . 5 [CO1]

**MATH-1051**

*[ Turn Over ]*

( 2 )

3. (a) Show that  $\frac{\pi^2}{9} < \int_{\pi/6}^{\pi/2} \frac{x}{\sin x} dx < \frac{2\pi^2}{9}$ .

(b) If  $f:[a,b] \rightarrow \mathbb{R}$  and  $\phi:[a,b] \rightarrow \mathbb{R}$  are both  $R$ -integrable on  $[a, b]$  and  $f$  is monotone decreasing on  $[a, b]$ , then show that  $\exists$  a point  $\xi$  in  $[a, b]$  such that

$$\int_a^b f(x)\phi(x)dx = f(a)\int_a^{\xi} \phi(x)dx + f(b)\int_{\xi}^b \phi(x)dx .$$

3+2=5 [CO1]

4. (a) If  $f:[a,b] \rightarrow \mathbb{R}$  is integrable on  $[a, b]$ , then show that the function  $F$  defined by  $F(x) = \int_a^x f(t)dt$ ,  $x \in [a, b]$  is continuous on  $[a, b]$ .

(b) Show that  $[x]$  is integrable on  $[0, 3]$  and evaluate  $\int_0^3 [x]dx$ .

3+2=5 [CO1]

5. (a) State Bonnet's form of 2nd M. V. T. in the integral calculus.

(b) The function  $f$  defined by

$$f(x) = \begin{cases} \frac{1}{n}, & \frac{1}{n+1} < x \leq \frac{1}{n} (n=1,2,3,\dots) . \\ 0, & x=0 \end{cases}$$

Show that  $f$  is  $R$ -integrable on  $[0, 1]$ . Evaluate  $\int_0^1 f dx$ .

1+4=5 [CO1]

6. Examine the convergence of  $\int_0^1 x^{m-1}(1-x)^{n-1} \log x dx$ .

5 [CO2]

7. (a) Find  $\int_0^1 \frac{dx}{(1-x^3)^{\frac{1}{3}}}$ .

(b) A function  $f$  is defined on  $[1, \infty)$  by  $f(x) = \frac{(-1)^{n-1}}{n}$  for  $n \leq x < n+1$  ( $n=1, 2, 3, \dots$ ). Examine the convergence of  $\int_1^\infty f(x)dx$ . 2+3=5 [CO2]

### PART—II (20 Marks)

1. Answer *any one* question :

4×1=4

(a) Using the definition of uniform convergence of sequence of functions, prove that the sequence of functions  $\{f_n\}$  defined by  $f_n(x) = \tan^{-1} nx$ ,  $\forall x \geq 0$ ,  $\forall n \in \mathbb{N}$  is uniformly convergent on  $[a, b]$  for all  $b > a > 0$  but the sequence of functions  $\{f_n\}$  is not uniformly convergent on  $[0, b]$  for any  $b > 0$ . 3+1=4 [CO1]

(b) Show that the sequence of functions  $\{f_n\}$  defined by  $f_n(x) = x^n$ ,  $x \in \mathbb{R}$  is point-wise convergent for  $x \in (-1, 1]$  and in this case find  $\lim_{n \rightarrow \infty} f_n(x)$ . Using the definition of uniform convergence of sequence of functions, prove that the above sequence of functions is uniformly convergent on  $[0, b]$  for all  $0 < b < 1$  but the sequence of functions  $\{f_n\}$  is not uniformly convergent on  $[0, 1]$ . 1+2+1=4 [CO1]

( 4 )

2. Answer *any two* questions :

4×2=8

(a) Let  $\{f_n\}$  be a sequence of functions defined on  $E(\subseteq R)$ .  
If  $\{f_n\}$  is point-wise convergent to  $f$  on  $E(\subseteq R)$  and  
 $M_n = \sup_{x \in E} |f_n(x) - f(x)|$  exists for all  $n \in \mathbb{N}$ , then  
 $\{f_n\}$  is uniformly convergent to  $f$  on  $E$  if and only if  
 $M_n \rightarrow 0$  as  $n \rightarrow \infty$ . 4 [CO2]

(b) Let  $\{f_n\}$  be a sequence of functions defined on  $E(\subseteq R)$ .  
If  $f_n(x) \leq M_n \forall x \in E$  and  $\forall n \in \mathbb{N}$  and  $\sum_{n=1}^{\infty} M_n$  is  
convergent, then  $\sum_{n=1}^{\infty} f_n(x)$  is uniformly convergent on  
 $E(\subseteq R)$ . 4 [CO2]

(c) Let  $R > 0$  be the radius of convergence of the power  
series  $\sum_{n=0}^{\infty} a_n x^n$ . Then prove that the radius of  
convergence of  $\sum_{n=1}^{\infty} n a_n x^{n-1}$  = the radius of  
convergence of  $\sum_{n=0}^{\infty} \frac{a_n}{n+1} x^{n+1} = R$ . 2+2=4 [CO2]

3. Answer *any one* question :

4×1=4

(a) Show that the series  $\sum_{n=1}^{\infty} f_n(x)$  converges uniformly on  
[0, 1], where

$$f_1(x) = \frac{x^2}{1+x}, f_n(x) = \frac{nx^2}{1+nx} - \frac{(n-1)x^2}{1+(n-1)x} \forall n-1 \in \mathbb{N}$$

4 [CO3]

( 5 )

- (b) Find the radius of convergence of the following power series : 4 [CO3]

$$\frac{1^2}{4^2} \frac{x}{3} + \frac{1^2 \cdot 5^2}{4^2 \cdot 8^2} \left(\frac{x}{3}\right)^2 + \frac{1^2 \cdot 5^2 \cdot 9^2}{4^2 \cdot 8^2 \cdot 12^2} \left(\frac{x}{3}\right)^3 + \frac{1^2 \cdot 5^2 \cdot 9^2 \cdot 13^2}{4^2 \cdot 8^2 \cdot 12^2 \cdot 16^2} \left(\frac{x}{3}\right)^4 + \dots$$

4. Answer *any one* question : 4×1=4

- (a) Assuming the power series expansion of  $(1+x^2)^{-1}$ , obtain the power series expansion of  $\tan^{-1}x$  and hence

obtain the sum of the series  $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{2n-1}$ .  
3+1=4 [CO4]

- (b) Express  $f(x) = x(\pi - x)$ ,  $0 < x < \pi$  as (i) Half-Range Fourier cosine series and (ii) Half-Range Fourier sine series by constructing two functions  $g(x)$  and  $h(x)$  of periodicity  $2\pi$ , where  $g(x)$  is even and  $h(x)$  is odd.

2+2=4 [CO4]

